

FERROELECTRIC PHASE TRANSITIONS IN A LATTICE PSEUDO-JAHN-TELLER MODEL

Dennis P. Clougherty

Department of Physics

University of Vermont

Burlington, VT 05403-0125

dpc@physics.uvm.edu

Keywords: ferroelectric, antiferroelectric, pseudo-Jahn-Teller

Abstract A lattice generalization of the $(b_{1u} \oplus b_{2g}) \otimes b_{3u}$ pseudo-Jahn-Teller model is analyzed within mean-field theory. Ferroelectric or antiferroelectric ground states are found for sufficiently large pseudo-Jahn-Teller coupling.

Introduction

Inspired by the family of fullerene-based polymeric structures synthesized [1, 2], a model describing vibronic effects in a linear chain of molecules is introduced and analyzed within mean-field theory. The zero-temperature phase diagrams at carrier concentrations $n = 1$ and 2 are presented for this model. Only a local $(b_{1u} \oplus b_{2g}) \otimes b_{3u}$ pseudo-Jahn-Teller interaction is considered here; spin and charge fluctuation effects are ignored.

The local symmetry of the molecule is taken to be D_{2h} for simplicity; the set of b -representations transform as components of a vector. The electronic structure of the molecule is taken to have a b_{1u} level as the HOMO and a b_{2g} level as the LUMO. A distortion of b_{3u} symmetry will vibronically couple these levels. For a molecule with some ionic bonding character, such a b_{3u} distortion produces an electric dipole moment in proportion to the displacement[3]. It is the ordering of these local dipole moments that is the concern of this work.

1. THE HAMILTONIAN

The Hamiltonian is given by

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_{\text{PJT}} + \mathcal{H}_{\text{elast}} \quad (1.1)$$

where

$$\mathcal{H}_0 = -t \sum_n (c_{1,n}^\dagger c_{1,n+1} + c_{2,n}^\dagger c_{2,n+1} + \text{h.c.}) + \sum_n \Delta c_{2,n}^\dagger c_{2,n} \quad (1.2)$$

where n runs over molecule sites, $c_{1,n}^\dagger$ ($c_{1,n}$) creates (annihilates) an electron in orbital b_{1u} at site n , $c_{2,n}^\dagger$ ($c_{2,n}$) creates (annihilates) an electron in orbital b_{2g} at site n , and Δ is the energy spacing between the b_{1u} and b_{2g} orbitals. The hopping matrix is taken to be diagonal in orbital type and spin, and the spin indices are suppressed for clarity.

The pseudo-Jahn-Teller Hamiltonian on the n th molecular site is

$$\mathcal{H}_{\text{PJT}} = -g \sum_n (c_{1,n}^\dagger c_{2,n} Q_n + \text{h.c.}) \quad (1.3)$$

where g is the pseudo-Jahn-Teller coupling strength, and Q_n is a displacement of b_{3u} symmetry on the n th molecule. The magnitude of the induced local electric dipole is proportional to Q_n . Hence, a ferrodistortive state is equated with a ferroelectric one.

$\mathcal{H}_{\text{elast}}$ is the elastic energy for the molecular displacements.

$$\mathcal{H}_{\text{elast}} = \frac{\kappa}{2} \sum_n Q_n^2 \quad (1.4)$$

The Hamiltonian can be written in k -space as

$$\mathcal{H} = \sum_{\alpha,k} \epsilon_\alpha(k) c_{\alpha,k}^\dagger c_{\alpha,k} - \frac{g\Delta}{\sqrt{N}} \sum_{k,q} (c_{1,k+q}^\dagger c_{2,k} Q_q + \text{h.c.}) + \frac{1}{2} \kappa \sum_q |Q_q|^2 \quad (1.5)$$

where $\epsilon_g(k) = \Delta - 2t \cos k$ and $\epsilon_u(k) = -2t \cos k$.

2. MEAN FIELD THEORY

With a single mode of distortion given by a non-vanishing Q_q , $\mathcal{H}$ can be diagonalized with a canonical transformation so that

$$\mathcal{H} = \sum_k (E_{k+} n_{k+} + E_{k-} n_{k-}) + \frac{\kappa}{2} |Q_q|^2 \quad (1.6)$$

where

$$E_{k\pm} = \frac{1}{2} (\epsilon_u(k) + \epsilon_g(k+q)) \pm \frac{1}{2} \sqrt{(\epsilon_g(k+q) - \epsilon_u(k))^2 + \frac{4g^2}{N} |Q_q|^2} \quad (1.7)$$

and $n_{k\pm}$ are the usual occupation number operators of the vibronic quasi-particles.

The ground state energy of a chain with n carriers ($0 \leq n \leq 2$) per molecule on average is obtained by fractionally filling the lowest energy vibronic band. In the continuum limit,

$$E(Q_q) = \frac{N}{\pi} \int_{-\pi n/2}^{\pi n/2} dk E_{k-} + \frac{\kappa}{2} |Q_q|^2 \quad (1.8)$$

Minimizing $E(Q_q)$ with respect to the distortion gives the mean-field equation to determine the magnitude of Q_q .

$$\frac{\pi\kappa}{2g^2} = \int_{-\pi n/2}^{\pi n/2} dk \frac{1}{\sqrt{\frac{4g^2}{N} |Q_q|^2 + (\Delta + 4t \sin q/2 \sin(k + q/2))^2}} \quad (1.9)$$

For sufficiently small coupling, this equation can not be satisfied. There is a q -dependent critical coupling strength g_c given by

$$g_c^2(q) = \frac{\frac{\pi\kappa\Delta}{2}}{\int_{-\pi n/2}^{\pi n/2} \frac{dk}{1 + u \sin q/2 \sin(k + q/2)}} \quad (1.10)$$

where $u = \frac{4t}{\Delta}$. For values of g above $g_c(q)$, a mean-field distortion of wavenumber q sets in; for values of g below $g_c(q)$, there is no such distortion.

3. THE PHASE DIAGRAM

For the case of an average occupancy of one carrier per molecule, the ground state is found to have a $q = 0$ distortion for $g > g_0 (= \sqrt{\frac{\kappa\Delta}{2}})$, consequently breaking inversion symmetry of the chain. The phase diagram is pictured in Fig. 1.

For the case of two carriers per molecule, the ground state has a $q = \pi$ distortion when

$$g^2 > \frac{g_0^2}{2} \sqrt{1 - u^2} \quad (1.11)$$

The results are summarized in Fig. 2.

Acknowledgments

This work was supported by the National Science Foundation through a grant for the Institute for Theoretical Atomic and Molecular Physics at Harvard University and the Smithsonian Astrophysical Observatory.




Figure 1 Zero-temperature phase diagram for $n = 1$.




Figure 2 Zero-temperature phase diagram for $n = 2$.

References

- [1] Pekker, S. *et al.*, *Science* **256**, 1077 (1994).
- [2] Stephens P. W. *et al.*, *Nature* **370**, 636 (1994).
- [3] Clougherty D. P. and Anderson F. G., *Phys. Rev. Lett.* **80**, 3735 (1998).